

CLAIMS

1. An arrangement for generating a representation of a feature in a surface defined by a mesh representation, the mesh comprising at a selected level a plurality of points including at least one point connected to a plurality of neighboring points by respective edges, the feature being defined in connection with the vertex and at least one of the neighboring points and the edge interconnecting the vertex and the at least one of the neighboring points in the mesh representation, the feature generating arrangement comprising:

- A. a weight vector generator module configured to generate at least one weight vector based on a parameterized subdivision rule defined at a plurality of levels, for which a value of at least one parameter differs at at least two levels in the mesh; and
- B. a feature representation generator module configured to use the at least one weight vector and positions of the vertex and the neighboring points to generate the representation of the feature.

2. An arrangement as defined in claim 1 in which weight vector generator module is configured to make use of values of the at least one parameter that differ at at least two levels are related by a selected mathematical function.

3. An arrangement as defined in claim 1 in which the feature is a smooth feature line.

4. An arrangement as defined in claim 3 in which the smooth feature line is defined in connection with the vertex and two neighboring points and edges interconnecting the vertex and the respective neighboring points, the weight vector generator module being configured to make use of parameterized subdivision rule having a parameter value associated with each of the edges along which the smooth feature line is defined.

5. An arrangement as defined in claim 4 in which the weight vector generator module is configured to make use of parameters associated with the edges along which the smooth feature line is defined whose values are the same.

6. An arrangement as defined in claim 5 in which the weight vector generator module is configured to make use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp crease along the edges along which the smooth feature line is defined, the values of the parameters being defined in the interval [0,1], where higher values define a sharper crease, the values of the parameters at a lower level being related to the values of the parameters at a higher level being related by

$$s(j+1) = (s(j))^2,$$

where $s(j)$ represents the values of the parameters at level "j" and $s(j+1)$ represents the values of the parameters at the higher level "j+1."

7. An arrangement as defined in claim 4 in which the weight vector generator module is configured to make use of parameters associated with the edges along which the smooth feature line is defined whose values differ.

8. An arrangement as defined in claim 7 in which the weight vector generator module is configured to make use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp crease along the edges along which the smooth feature line is defined, the values of the parameters being defined in the interval [0,1], where higher values define a sharper crease, the values of the parameters at a lower level being related to the values of the parameters at a higher level being related by

$$s_1(j+1) = \left(\frac{3}{4}s_1(j) + \frac{1}{4}s_2(j) \right)^2,$$

and

$$s_2(j+1) = \left(\frac{1}{4}s_1(j) + \frac{3}{4}s_2(j) \right)^2,$$

where $s_1(j)$ and $s_2(j)$ represent the values of the parameters associated with the respective edges at level "j," and $s_1(j+1)$ and $s_2(j+1)$ represent the values of the parameters associated with the respective edges at the higher level "j+1."

9. An arrangement as defined in claim 4 in which the mesh comprises a triangular mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1, \dots, K$ are at respective positions $c^j(k)$, and in which the weight vector generator module is configured to make use of a parameterized subdivision rule $S_{sc,T,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j,$$

where subdivision rule $S_{sc,T,K,L}$ is given by

10062192-020102

A1
CMT

$$(S_{sc,T,K,L}(s_1, s_2))_{l,m} = \left\{ \begin{array}{ll} (1-s_3)(1-a(K)) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ (1-s_3)\frac{a(K)}{K} + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ (1-s_3)\frac{a(K)}{K} & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ \frac{3}{8} + \frac{1}{8}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{3}{8} + \frac{1}{8}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{8}(1-s_2) & \text{if } l=1, m=2 \text{ or } K \\ \frac{1}{8}(1-s_1) & \text{if } l=L+1, m=L \text{ or } L+2 \\ \frac{3}{8} & \\ \frac{1}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{8} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=L+2, \dots, K, m=l+1 \\ & \text{or } l=K, m=1 \\ 0 & \text{otherwise} \end{array} \right.$$

8

9

10

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point

11 $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex

12 $v_q(0)$, $s_3 = \frac{1}{2}(s_1 + s_2)$, and

$$a(K) = \frac{5}{8} - \left(\frac{3 + 2 \cos\left(\frac{2\pi}{K}\right)}{8} \right)^2$$

10. An arrangement as defined in claim 9 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator module being configured to determine a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^K \left(l_{LP}(s_1, s_2) \right)_i c^j(i)$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc,T,K,L,LP}(s_1, s_2)$$

where

$$S_{sc,T,K,L,LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc,T,K,L}(s_1(j), s_2(j))$$

where $S_{sc,T,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc,T,K,L,LP}(s_1, s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \left(\frac{\omega(K)}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \dots, \frac{1}{\varpi(K) + K} \right),$$

where

$$\varpi(K) = \frac{3K}{8a(K)}$$

11. An arrangement as defined in claim 10 in which the weight vector generator module is configured to generate an approximation for the limit point weight vector l_{LP} using a polynomial approximation methodology.

12. An arrangement as defined in claim 11 in which the weight vector generator module is configured to generate the approximation for the limit point weight vector l_{LP} in accordance with the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2,$$

in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_2^2 + b_{i5}s_1s_2,$$

in an asymmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

13. An arrangement as defined in claim 12 in which weight vector generator module is configured to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

14. An arrangement as defined in claim 10 in which the weight vector generator module is configured to generate an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

15. An arrangement as defined in claim 14 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

16. An arrangement as defined in claim 15 in which the weight vector generator module is configured to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J),$$

where "J" is a predetermined integer and where

$$(l_{LP}(J)(s_1, s_2))_m = (S_{sc, T, K, L, LP}(J)(s_1, s_2))_{1, m},$$

7 where

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)) ,$$

8

9 with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1} ,$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and

where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

13

1 17. An arrangement as defined in claim 9 in which the representation of the feature is defined by a
2 tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the
3 feature representation generator module being configured to determine the tangent vector $e_c(q)$ in
4 accordance with

$$e_c(q) = \sum_{i=0}^K (l_c(s_1, s_2))_i c^j(i)$$

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c = \lim_{j \rightarrow \infty} \frac{l_c(J)}{\|l_c(J)\|}$$

where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, and where

$$l_c(J) = (0, 1, 0, \dots, -1, 0, \dots) \cdot \prod_{j=J}^{j_D} S_{sc, T, K, L}(s_1(j), s_2(j))$$

where two non-zero components of the row vector on the right hand side are a "one" at position "one" in the row vector, and a "negative one" at position $\frac{K}{2} + 1$, and where $S_{sc, T, K, L}(s_1(j), s_2(j))$ corresponds to $S_{sc, T, K, L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product.

18. An arrangement as defined in claim 17 in which the weight vector generator module is configured to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

19. An arrangement as defined in claim 18 in which the weight vector generator module is configured to generate the approximation for the tangent vector weight vector l_c in accordance with the polynomial

$$(l_c)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2) ,$$

in the anti-symmetric case, or the polynomial

$$(l_c)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3 ,$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

20. An arrangement as defined in claim 19 in which weight vector generator module is configured to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right) ,$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

21. An arrangement as defined in claim 17 in which the weight vector generator module is configured to generate an approximation for the tangent vector weight vector l_c using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

22. An arrangement as defined in claim 21 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

23. An arrangement as defined in claim 22 in which the weight vector generator module is configured to generate the approximation for the tangent vector weight vector l_c in accordance with

$$l_c \approx \sum_{J=0}^3 b_J l_c(J) ,$$

where "J" is a predetermined integer and where

$$l_c(J)(s_1, s_2) = d(K)^J v_c \cdot S_{sc,T,K,L,LP}(J)(s_1, s_2) ,$$

where vector v_c is given by

$$v_c = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right) ,$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}} ,$$

and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)) ,$$

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1},$$

13

14 where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

24. An arrangement as defined in claim 9 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being across the smooth feature line, the feature representation generator module being configured to determine the tangent vector $e_c(q)$ in accordance with

$$e_s(q) = \sum_{i=0}^K (l_s(s_1, s_2))_i c^j(i)$$

where $l_s(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s = \lim_{j \rightarrow \infty} \frac{l_s(J)}{\|l_s(J)\|}$$

8 where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, where

$$l_s(J) = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right) \cdot \prod_{j=J}^{J_D} S_{sc,T,K,L}(s_1(j), s_2(j)) \quad ,$$

9
10 where $S_{sc,T,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th"
11 level in the matrix product.

12 25. An arrangement as defined in claim 24 in which the weight vector generator module is
13 configured to generate the tangent vector weight vector l_c using a polynomial approximation
methodology.

14 26. An arrangement as defined in claim 25 in which the weight vector generator module is
15 configured to generate the approximation for the tangent vector weight vector l_s in accordance with
16 the polynomial

$$(l_s)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2) \quad ,$$

17 4
18 5 in the anti-symmetric case ($i = 0$ or $\frac{K}{4} + 1$), or the polynomial

$$(l_s)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3 \quad ,$$

19 6
20 7 in the non-symmetric case ("i" otherwise), in which the coefficients b_{ij} are determined by a least
21 8 squares methodology and the values of the parameters s_1 and s_2 are at selected values.

27. An arrangement as defined in claim 26 in which weight vector generator module is configured to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

28. An arrangement as defined in claim 23 in which the weight vector generator module is configured to generate an approximation for the tangent vector weight vector l_s using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

29. An arrangement as defined in claim 28 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

30. An arrangement as defined in claim 29 in which the weight vector generator module is configured to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_s \approx \sum_{J=0}^3 b_J l_{s_J}(J),$$

where "J" is a predetermined integer and where

$$I_S(J)(s_1, s_2) = d(K)^J v_S \cdot S_{sc,T,K,L,LP}(J)(s_1, s_2),$$

where vector v_S is given by

$$v_S = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right),$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}},$$

and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)),$$

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1},$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

31. An arrangement as defined in claim 4 in which the mesh comprises a quadrilateral mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1, \dots, 2K$ are at respective positions $c^j(k)$, and in which the weight vector generator module is configured to make use of a parameterized subdivision rule $S_{sc,Q,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j,$$

where subdivision rule $S_{sc,Q,K,L}$ is given by

10062192-020102

A1
Cmt

$$(S_{sc,Q,K,L}(s_1, s_2))_{l,m} = \begin{cases} \left(1-s_3\right)\left(1-\frac{7}{4K}\right) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ \left(1-s_3\right)\left(\frac{3}{2K^2}\right) + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ \left(1-s_3\right)\left(\frac{3}{2K^2}\right) & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ \left(1-s_3\right)\left(\frac{1}{4K^2}\right) & \text{if } l=0, m=K+1, \dots, 2K \\ \frac{3}{8}(1-s_2) + \frac{1}{2}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{1}{16}(1-s_2) & \text{if } l=L+1, m=2, K, \\ & K+1 \text{ or } 2K \\ \frac{3}{8}(1-s_1) + \frac{1}{2}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{16}(1-s_1) & \text{if } l=L+1, m=L, L+2, \\ & K+1 \text{ or } K+L+1 \\ \frac{3}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{16} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=2, \dots, L, m=K+l-1 \\ & \text{or } l=2, \dots, L, m=K+l \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=L+2, \dots, K-1, m=l+1 \\ & \text{or } l=K, m=1 \\ & \text{or } l=L+2, \dots, K, m=K+l-1 \\ & \text{or } l=L+2, \dots, K, m=K+l \\ \frac{1}{4} & \text{if } l=K+1, \dots, 2K-1, m=0, \\ & l-K, l-K+1 \text{ or } l \\ & \text{or } l=2K, m=0, K, 1 \text{ or } 2K \\ 0 & \text{otherwise} \end{cases}$$

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq 2K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex $v_q(0)$, and $s_3 = \frac{1}{2}(s_1 + s_2)$.

32. An arrangement as defined in claim 31 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator module being configured to determine a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^{2K} (l_{LP}(s_1, s_2))_i c^j(i) ,$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc,Q,K,L,LP}(s_1, s_2) ,$$

where

$$S_{sc,Q,K,L,LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc,Q,K,L}(s_1(j), s_2(j)) ,$$

where $S_{sc,Q,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,Q,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc,Q,K,L,LP}(s_1, s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \frac{1}{K(K+5)} (K^2, 4, \dots, 4, 1, \dots, 1) .$$

13

1 33. An arrangement as defined in claim 32 in which the weight vector generator module is
2 configured to generate an approximation for the limit point weight vector l_{LP} using a polynomial
3 approximation methodology.

1 34. An arrangement as defined in claim 33 in which the weight vector generator module is
2 configured to generate the approximation for the limit point weight vector l_{LP} in accordance with the
3 polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2, \quad ,$$

5 in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2, \quad ,$$

6
7 in an asymmetric case, in which the coefficients b_{ij} are determined by a least squares methodology
8 and the values of the parameters s_1 and s_2 are at selected values.

1 35. An arrangement as defined in claim 34 in which weight vector generator module is configured
2 to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right), \quad ,$$

3

4 where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

36. An arrangement as defined in claim 32 in which the weight vector generator module is configured to generate an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

37. An arrangement as defined in claim 36 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

38. An arrangement as defined in claim 37 in which the weight vector generator module is configured to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J),$$

where "J" is a predetermined integer and where

$$(l_{LP}(J)(s_1, s_2))_m = (S_{sc,Q,K,L,LP}(J)(s_1, s_2))_{1,m},$$

where

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j)),$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1},$$

10

11 where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

12

1

39. An arrangement as defined in claim 31 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the feature representation generator module being configured to determine the tangent vector $e_c(q)$ in accordance with

4

$$e_c(q) = \sum_{i=0}^{2K} \left(l_c(s_1, s_2) \right)_i c^j(i) ,$$

5

6 where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c(s_1, s_2) = d(K)^K v_c \cdot S_{sc,Q,K,L,LP}(s_1(j), s_2(j)) ,$$

7

8 where

$$S_{sc,Q,K,L,LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc,Q,K,L}(s_1(j), s_2(j)) ,$$

and vector v_c is defined as

$$(v_c)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \cos \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \cos \frac{2\pi(i-K-1)}{K} + \cos \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases}$$

where

$$A_k = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)}$$

and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_k}{16} + \frac{1}{4}}$$

40. An arrangement as defined in claim 39 in which the weight vector generator module is configured to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

41. An arrangement as defined in claim 40 in which the weight vector generator module is configured to generate the approximation for the tangent vector weight vector l_c in accordance with the polynomial

$$(l_c)_i \approx b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2)$$

in the anti-symmetric case, or the polynomial

$$(l_c)_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

42. An arrangement as defined in claim 41 in which weight vector generator module is configured to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right)$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

43. An arrangement as defined in claim 39 in which the weight vector generator module is configured to generate an approximation for the tangent vector weight vector l_c using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

44. An arrangement as defined in claim 43 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

45. An arrangement as defined in claim 44 in which the weight vector generator module is configured to generate the approximation for the tangent vector weight vector l_c in accordance with

$$l_c \approx \sum_{J=0}^3 b_J l_c(J) ,$$

where "J" is a predetermined integer and where

$$l_c(J)(s_1, s_2) = d(K)^J v_c \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2) ,$$

where vector v_c is given by

$$v_c = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right) ,$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}} ,$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j)) ,$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1} ,$$

14 where I_{K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

1 46. An arrangement as defined in claim 31 in which the representation of the feature is defined by
 2 a tangent vector associated with the vertex, the tangent vector being across the smooth feature line,
 3 the feature representation generator module being configured to determine the tangent vector $e_c(q)$
 4 in accordance with

$$e_s(q) = \sum_{i=0}^{2K} (l_s(s_1, s_2))_i c^j(i) ,$$

5
 6 where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s(s_1, s_2) = d(K)^K v_s \cdot S_{sc, Q, K, L, LP}(s_1(j), s_2(j)) ,$$

7
 8 where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

and vector v_c is defined as

$$(v_s)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \sin \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \sin \frac{2\pi(i-K-1)}{K} + \sin \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases},$$

where

$$A_K = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)},$$

and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_K}{16} + \frac{1}{4}}.$$

47. An arrangement as defined in claim 46 in which the weight vector generator module is configured to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

48. An arrangement as defined in claim 47 in which the weight vector generator module is configured to generate the approximation for the tangent vector weight vector l_s in accordance with the polynomial

$$(l_s)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2),$$

in the anti-symmetric case, or the polynomial

$$(l_s)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

49. An arrangement as defined in claim 48 in which weight vector generator module is configured to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right)$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

50. An arrangement as defined in claim 46 in which the weight vector generator module is configured to generate an approximation for the tangent vector weight vector l_s using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

51. An arrangement as defined in claim 50 in which the weight vector generator module is configured to perform the extrapolation approximation methodology such that $M=3$.

52. An arrangement as defined in claim 51 in which the weight vector generator module is configured to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_s \approx \sum_{J=0}^3 b_J l_s(J)$$

where "J" is a predetermined integer and where

$$l_s(J)(s_1, s_2) = d(K)^J v_s \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2)$$

where vector v_s is given by

$$v_s = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right)$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}}$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j))$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1}$$

14

15 where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

16

1

2

3

4

5

6

7

8

9

10

11

12

13

53. A computer program product for use in connection with a computer to provide an arrangement for generating a representation of a feature in a surface defined by a mesh representation, the mesh comprising at a selected level a plurality of points including at least one point connected to a plurality of neighboring points by respective edges, the feature being defined in connection with the vertex and at least one of the neighboring points and the edge interconnecting the vertex and the at least one of the neighboring points in the mesh representation, the computer program product comprising:

- A. a weight vector generator module configured to enable the computer to generate at least one weight vector based on a parameterized subdivision rule defined at a plurality of levels, for which a value of at least one parameter differs at at least two levels in the mesh; and
- B. a feature representation generator module configured to enable the computer to use the at least one weight vector and positions of the vertex and the neighboring points to generate the representation of the feature.

54. A computer program product as defined in claim 53 in which weight vector generator module is configured to enable the computer to make use of values of the at least one parameter that differ at at least two levels are related by a selected mathematical function.

55. A computer program product as defined in claim 53 in which the feature is a smooth feature line.

56. A computer program product as defined in claim 55 in which the smooth feature line is defined in connection with the vertex and two neighboring points and edges interconnecting the vertex and the respective neighboring points, the weight vector generator module being configured to enable the computer to make use of parameterized subdivision rule having a parameter value associated with each of the edges along which the smooth feature line is defined.

57. A computer program product as defined in claim 56 in which the weight vector generator module is configured to enable the computer to make use of parameters associated with the edges along which the smooth feature line is defined whose values are the same.

58. A computer program product as defined in claim 57 in which the weight vector generator module is configured to enable the computer to make use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp crease along the edges along which the smooth feature line is defined, the values of the parameters being defined in the interval [0,1], where higher values define a sharper crease, the values of the parameters at a lower level being related to the values of the parameters at a higher level being related by

$$s(j+1) = (s(j))^2,$$

where $s(j)$ represents the values of the parameters at level "j" and $s(j+1)$ represents the values of the parameters at the higher level "j+1."

59. A computer program product as defined in claim 56 in which the weight vector generator module is configured to enable the computer to make use of parameters associated with the edges along which the smooth feature line is defined whose values differ.

60. A computer program product as defined in claim 59 in which the weight vector generator module is configured to enable the computer to make use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp crease along the edges along which the smooth feature line is defined, the values of the parameters being defined in the interval [0,1], where higher values define a sharper crease, the values of the parameters at a lower level being related to the values of the parameters at a higher level being related by

$$s_1(j+1) = \left(\frac{3}{4}s_1(j) + \frac{1}{4}s_2(j) \right)^2,$$

and

$$s_2(j+1) = \left(\frac{1}{4}s_1(j) + \frac{3}{4}s_2(j) \right)^2,$$

where $s_1(j)$ and $s_2(j)$ represent the values of the parameters associated with the respective edges at level "j," and $s_1(j+1)$ and $s_2(j+1)$ represent the values of the parameters associated with the respective edges at the higher level "j+1."

61. A computer program product as defined in claim 56 in which the mesh comprises a triangular mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1,...,K$ are at respective positions $c^j(k)$, and in which the weight vector generator module is configured to enable the computer to make use of a parameterized subdivision rule $S_{sc,T,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j$$

7

8 where subdivision rule $S_{sc,T,K,L}$ is given by

$$(S_{sc,T,K,L}(s_1, s_2))_{l,m} = \begin{cases} (1-s_3)(1-a(K)) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ (1-s_3)\frac{a(K)}{K} + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ (1-s_3)\frac{a(K)}{K} & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ \frac{3}{8} + \frac{1}{8}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{3}{8} + \frac{1}{8}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{8}(1-s_2) & \text{if } l=1, m=2 \text{ or } K \\ \frac{1}{8}(1-s_1) & \text{if } l=L+1, m=L \text{ or } L+2 \\ \frac{3}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{8} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=L+2, \dots, K, m=l+1 \\ & \text{or } l=K, m=1 \\ 0 & \text{otherwise} \end{cases}$$

10062192-020102

A/CMT

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex $v_q(0)$, $s_3 = \frac{1}{2}(s_1 + s_2)$, and

$$a(K) = \frac{5}{8} - \left(\frac{3 + 2 \cos\left(\frac{2\pi}{K}\right)}{8} \right)^2$$

62. A computer program product as defined in claim 61 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator module being configured to enable the computer to determine a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^K \left(l_{LP}(s_1, s_2) \right)_i c^j(i),$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc,T,K,L,LP}(s_1, s_2),$$

where

$$S_{sc,T,K,L,LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc,T,K,L}(s_1(j), s_2(j)),$$

where $S_{sc,T,K,L}(s_1(j),s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc,T,K,L,LP}(s_1,s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \left(\frac{\omega(K)}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \dots, \frac{1}{\varpi(K) + K} \right),$$

where

$$\omega(K) = \frac{3K}{8a(K)}$$

63. A computer program product as defined in claim 62 in which the weight vector generator module is configured to enable the computer to generate an approximation for the limit point weight vector l_{LP} using a polynomial approximation methodology.

64. A computer program product as defined in claim 63 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector l_{LP} in accordance with the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2,$$

in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_2^2 + b_{i5}s_1s_2,$$

in an asymmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

65. A computer program product as defined in claim 64 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

66. A computer program product as defined in claim 62 in which the weight vector generator module is configured to enable the computer to generate an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

67. A computer program product as defined in claim 66 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

68. A computer program product as defined in claim 67 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J)$$

4

5 where "J" is a predetermined integer and where

$$\left(l_{LP}(J)(s_1, s_2) \right)_m = \left(S_{sc,T,K,L,LP}(J)(s_1, s_2) \right)_{1,m},$$

6

7 where

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)),$$

8

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1},$$

10

11 where I_{K+1} is the "K+1" by "K+1" identity matrix, and

12

where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

69. A computer program product as defined in claim 61 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the feature representation generator module being configured to enable the computer to determine the tangent vector $e_c(q)$ in accordance with

$$e_c(q) = \sum_{i=0}^K (l_c(s_1, s_2))_i c^j(i) ,$$

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c = \lim_{j \rightarrow \infty} \frac{l_c(J)}{\|l_c(J)\|} ,$$

where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, and where

$$l_C(J) = (0, 1, 0, \dots, -1, 0, \dots) \cdot \prod_{j=J}^{j_D} S_{sc,T,K,L}(s_1(j), s_2(j)) ,$$

where two non-zero components of the row vector on the right hand side are a "one" at position "one" in the row vector, and a "negative one" at position $\frac{K}{2} + 1$, and where $S_{sc,T,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product.

70. A computer program product as defined in claim 69 in which the weight vector generator module is configured to enable the computer to generate the tangent vector weight vector l_C using a polynomial approximation methodology.

71. A computer program product as defined in claim 70 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_C in accordance with the polynomial

$$(l_C)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2) ,$$

in the anti-symmetric case, or the polynomial

$$(l_C)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3 ,$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

72. A computer program product as defined in claim 71 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

73. A computer program product as defined in claim 69 in which the weight vector generator module is configured to enable the computer to generate an approximation for the tangent vector weight vector l_c using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

74. A computer program product as defined in claim 73 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

75. A computer program product as defined in claim 74 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_c in accordance with

$$l_c \approx \sum_{J=0}^3 b_J l_c(J),$$

where "J" is a predetermined integer and where

$$l_c(J)(s_1, s_2) = d(K)^J v_c \cdot S_{sc,T,K,L,LP}(J)(s_1, s_2),$$

where vector v_c is given by

$$v_c = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right),$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}},$$

and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)),$$

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1},$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

76. A computer program product as defined in claim 61 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being across the smooth feature line, the feature representation generator module being configured to enable the computer to determine the tangent vector $e_c(q)$ in accordance with

$$e_s(q) = \sum_{i=0}^K (l_s(s_1, s_2))_i c^j(i)$$

where $l_s(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s = \lim_{j \rightarrow \infty} \frac{l_s(j)}{\|l_s(j)\|}$$

where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, where

$$l_s(j) = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right) \cdot \prod_{j=J}^{j_D} S_{sc,T,K,L}(s_1(j), s_2(j)) \quad ,$$

where $S_{sc,T,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product.

77. A computer program product as defined in claim 76 in which the weight vector generator module is configured to enable the computer to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

78. A computer program product as defined in claim 77 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_s in accordance with the polynomial

$$(l_s)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2) \quad ,$$

in the anti-symmetric case, or the polynomial

$$(l_s)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3 \quad ,$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

79. A computer program product as defined in claim 78 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

80. A computer program product as defined in claim 75 in which the weight vector generator module is configured to enable the computer to generate an approximation for the tangent vector weight vector l_s using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

81. A computer program product as defined in claim 80 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

82. A computer program product as defined in claim 81 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_s \approx \sum_{J=0}^3 b_J l_s(J)$$

where "J" is a predetermined integer and where

$$l_s(J)(s_1, s_2) = d(K)^J v_s \cdot S_{sc, T, K, L, LP}(J)(s_1, s_2)$$

7 where vector v_s is given by

$$v_s = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right),$$

8

9 dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}}$$

and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)),$$

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) = I_{K+1},$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

A1
Gut

83. A computer program product as defined in claim 56 in which the mesh comprises a quadrilateral mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1, \dots, 2K$ are at respective positions $c^j(k)$, and in which the weight vector generator module is configured to enable the computer to make use of a parameterized subdivision rule $S_{sc,Q,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j,$$

where subdivision rule $S_{sc,Q,K,L}$ is given by

A1 cut

$$(S_{sc,Q,K,L}(s_1, s_2))_{l,m} = \begin{cases} (1-s_3)\left(1-\frac{7}{4K}\right) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ (1-s_3)\left(\frac{3}{2K^2}\right) + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ (1-s_3)\left(\frac{3}{2K^2}\right) & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ (1-s_3)\left(\frac{1}{4K^2}\right) & \text{if } l=0, m=K+1, \dots, 2K \\ \frac{3}{8}(1-s_2) + \frac{1}{2}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{1}{16}(1-s_2) & \text{if } l=L+1, m=2, K, \\ & K+1 \text{ or } 2K \\ \frac{3}{8}(1-s_1) + \frac{1}{2}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{16}(1-s_1) & \text{if } l=L+1, m=L, L+2, \\ & K+1 \text{ or } K+L+1 \\ \frac{3}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{16} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=2, \dots, L, m=K+l-1 \\ & \text{or } l=2, \dots, L, m=K+l \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=L+2, \dots, K-1, m=l+1 \\ & \text{or } l=K, m=1 \\ & \text{or } l=L+2, \dots, K, m=K+l-1 \\ & \text{or } l=L+2, \dots, K, m=K+l \\ \frac{1}{4} & \text{if } l=K+1, \dots, 2K-1, m=0, \\ & l-K, l-K+1 \text{ or } l \\ & \text{or } l=2K, m=0, K, 1 \text{ or } 2K \\ 0 & \text{otherwise} \end{cases}$$

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq 2K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex $v_q(0)$, and $s_3 = \frac{1}{2}(s_1 + s_2)$.

84. A computer program product as defined in claim 83 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator module being configured to enable the computer to determine a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^{2K} (l_{LP}(s_1, s_2))_i c^j(i),$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc,Q,K,L,LP}(s_1, s_2),$$

where

$$S_{sc,Q,K,L,LP}(s_1, s_2) = \prod_{j=0}^{j_D} S_{sc,Q,K,L}(s_1(j), s_2(j)),$$

where $S_{sc,Q,K,L}(s_1(j), s_2(j))$ corresponds to $S_{sc,Q,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc,Q,K,L,LP}(s_1, s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \frac{1}{K(K+5)} (K^2, 4, \dots, 4, 1, \dots, 1)$$

85. A computer program product as defined in claim 84 in which the weight vector generator module is configured to enable the computer to generate an approximation for the limit point weight vector l_{LP} using a polynomial approximation methodology.

86. A computer program product as defined in claim 85 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector l_{LP} in accordance with the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2,$$

in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2,$$

in an asymmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

87. A computer program product as defined in claim 86 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

88. A computer program product as defined in claim 84 in which the weight vector generator module is configured to enable the computer to generate an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

89. A computer program product as defined in claim 88 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

90. A computer program product as defined in claim 89 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J) ,$$

where "J" is a predetermined integer and where

$$(l_{LP}(J)(s_1, s_2))_m = (S_{sc, Q, K, L, LP}(J)(s_1, s_2))_{1, m} ,$$

where

$$S_{sc, Q, K, L, LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

with

$$S_{sc, Q, K, L, LP}(0)(s_1, s_2) := I_{2K+1} ,$$

10

11 where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

AI
CMT

10052192200102

1

2

3

4

91. A computer program product as defined in claim 83 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the feature representation generator module being configured to enable the computer to determine the tangent vector $e_c(q)$ in accordance with

$$e_c(q) = \sum_{i=0}^{2K} (l_c(s_1, s_2))_i c^j(i) ,$$

5

6

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c(s_1, s_2) = d(K)^K v_c \cdot S_{sc, Q, K, L, LP}(s_1(j), s_2(j)) ,$$

7

8

where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

and vector v_c is defined as

$$(v_c)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \cos \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \cos \frac{2\pi(i-K-1)}{K} + \cos \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases}$$

where

$$A_k = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)}$$

and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_k}{16} + \frac{1}{4}}$$

92. A computer program product as defined in claim 91 in which the weight vector generator module is configured to enable the computer to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

93. A computer program product as defined in claim 92 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_c in accordance with the polynomial

$$(l_c)_i \approx b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2)$$

in the anti-symmetric case, or the polynomial

$$(l_C)_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

94. A computer program product as defined in claim 93 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right)$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

95. A computer program product as defined in claim 91 in which the weight vector generator module is configured to enable the computer to generate an approximation for the tangent vector weight vector l_C using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

96. A computer program product as defined in claim 95 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

97. A computer program product as defined in claim 96 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_C in accordance with

$$l_C \approx \sum_{J=0}^3 b_J l_C(J) ,$$

where "J" is a predetermined integer and where

$$l_C(J)(s_1, s_2) = d(K)^J v_C \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2) ,$$

where vector v_C is given by

$$v_C = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right) ,$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}} ,$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j)) ,$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1} ,$$

14

15

where I_{K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

cut

6

1

2

3

4

98. A computer program product as defined in claim 83 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being across the smooth feature line, the feature representation generator module being configured to enable the computer to determine the tangent vector $e_c(q)$ in accordance with

$$e_s(q) = \sum_{i=0}^{2K} (l_s(s_1, s_2))_i c'(i) ,$$

5

6

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s(s_1, s_2) = d(K)^K v_s \cdot S_{sc, Q, K, L, LP}(s_1(j), s_2(j)) ,$$

7

8

where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

and vector v_c is defined as

$$(v_s)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \sin \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \sin \frac{2\pi(i-K-1)}{K} + \sin \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases},$$

where

$$A_K = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)},$$

and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_K}{16} + \frac{1}{4}}$$

99. A computer program product as defined in claim 98 in which the weight vector generator module is configured to enable the computer to generate the tangent vector weight vector l_c using a polynomial approximation methodology.

100. A computer program product as defined in claim 99 in which the weight vector generator module is configured to enable the computer to generate the approximation for the tangent vector weight vector l_s in accordance with the polynomial

$$(l_s)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2),$$

in the anti-symmetric case, or the polynomial

$$(l_s)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3,$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

101. A computer program product as defined in claim 100 in which weight vector generator module is configured to enable the computer to select values of s_1 and s_2 in accordance with

A1
cont

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

102. A computer program product as defined in claim 98 in which the weight vector generator module is configured to enable the computer to generate an approximation for the tangent vector weight vector l_s using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

103. A computer program product as defined in claim 102 in which the weight vector generator module is configured to enable the computer to perform the extrapolation approximation methodology such that $M=3$.

104. A computer program product as defined in claim 103 in which the weight vector generator module is configured to enable the computer to generate the approximation for the limit point weight vector weight vector in accordance with

$$l_s \approx \sum_{J=0}^3 b_J l_s(J) ,$$

where "J" is a predetermined integer and where

$$l_s(J)(s_1, s_2) = d(K)^J v_s \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2) ,$$

where vector v_s is given by

$$v_s = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right) ,$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}} ,$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j)) ,$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1} ,$$

14

15 where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

AI
Cont

16

105. A method for generating a representation of a feature in a surface defined by a mesh representation, the mesh comprising at a selected level a plurality of points including at least one point connected to a plurality of neighboring points by respective edges, the feature being defined in connection with the vertex and at least one of the neighboring points and the edge interconnecting the vertex and the at least one of the neighboring points in the mesh representation, the method comprising:

- A. a weight vector generator step configured to generate at least one weight vector based on a parameterized subdivision rule defined at a plurality of levels, for which a value of at least one parameter differs at at least two levels in the mesh, and
- B. a feature representation generator step configured to use the at least one weight vector and positions of the vertex and the neighboring points to generate the representation of the feature.

1 106. A method as defined in claim 105 in which weight vector generator step includes the step of
2 making use of values of the at least one parameter that differ at at least two levels are related by a
3 selected mathematical function.

1 107. A method as defined in claim 105 in which the feature is a smooth feature line.

1 108. A method as defined in claim 107 in which the smooth feature line is defined in connection
2 with the vertex and two neighboring points and edges interconnecting the vertex and the respective
3 neighboring points, the weight vector generator step includes the step of making use of
4 parameterized subdivision rule having a parameter value associated with each of the edges along
5 which the smooth feature line is defined.

1 109. A method as defined in claim 108 in which the weight vector generator step includes the step
2 of making use of parameters associated with the edges along which the smooth feature line is defined
3 whose values are the same.

1 110. A method as defined in claim 109 in which the weight vector generator step includes the step
2 of making use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp
3 crease along the edges along which the smooth feature line is defined, the values of the parameters
4 being defined in the interval [0,1], where higher values define a sharper crease, the values of the
5 parameters at a lower level being related to the values of the parameters at a higher level being
6 related by

$$s(j+1) = (s(j))^2,$$

7
8 where $s(j)$ represents the values of the parameters at level "j" and $s(j+1)$ represents the values of the
9 parameters at the higher level "j+1."

111. A method as defined in claim 108 in which the weight vector generator step includes the step of making use of parameters associated with the edges along which the smooth feature line is defined whose values differ.

112. A method as defined in claim 111 in which the weight vector generator step includes the step of making use of the parameters that are in relation to a subdivision rule that, in turn, reflects a sharp crease along the edges along which the smooth feature line is defined, the values of the parameters being defined in the interval [0,1], where higher values define a sharper crease, the values of the parameters at a lower level being related to the values of the parameters at a higher level being related by

$$s_1(j+1) = \left(\frac{3}{4}s_1(j) + \frac{1}{4}s_2(j) \right)^2,$$

and

$$s_2(j+1) = \left(\frac{1}{4}s_1(j) + \frac{3}{4}s_2(j) \right)^2,$$

where $s_1(j)$ and $s_2(j)$ represent the values of the parameters associated with the respective edges at level "j," and $s_1(j+1)$ and $s_2(j+1)$ represent the values of the parameters associated with the respective edges at the higher level "j+1."

113. A method as defined in claim 108 in which the mesh comprises a triangular mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1,...,K$ are at respective positions $c^j(k)$, and in which the weight vector generator step includes the step of making use of a parameterized subdivision rule $S_{sc,T,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j$$

6

7

where subdivision rule $S_{sc,T,K,L}$ is given by

$$(S_{sc,T,K,L}(s_1, s_2))_{l,m} = \begin{cases} (1-s_3)(1-a(K)) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ (1-s_3)\frac{a(K)}{K} + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ (1-s_3)\frac{a(K)}{K} & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ \frac{3}{8} + \frac{1}{8}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{3}{8} + \frac{1}{8}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{8}(1-s_2) & \text{if } l=1, m=2 \text{ or } K \\ \frac{1}{8}(1-s_1) & \text{if } l=L+1, m=L \text{ or } L+2 \\ \frac{3}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{8} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=L+2, \dots, K, m=l+1 \\ & \text{or } l=K, m=1 \\ 0 & \text{otherwise} \end{cases}$$

A1
CMT

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex $v_q(0)$, $s_3 = \frac{1}{2}(s_1 + s_2)$, and

$$a(K) = \frac{5}{8} - \left(\frac{3 + 2 \cos\left(\frac{2\pi}{K}\right)}{8} \right)^2$$

114. A method as defined in claim 113 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator step includes the step of determining a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^K \left(l_{LP}(s_1, s_2) \right)_i c^j(i)$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc,T,K,L,LP}(s_1, s_2)$$

where

$$S_{sc,T,K,L,LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc,T,K,L}(s_1(j), s_2(j))$$

where $S_{sc,T,K,L}(s_1(j),s_2(j))$ corresponds to $S_{sc,T,K,L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc,T,K,L,LP}(s_1,s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \left(\frac{\omega(K)}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \frac{1}{\varpi(K) + K}, \dots, \frac{1}{\varpi(K) + K} \right),$$

where

$$\varpi(K) = \frac{3K}{8a(K)}$$

115. A method as defined in claim 114 in which the weight vector generator step includes the step of generating an approximation for the limit point weight vector l_{LP} using a polynomial approximation methodology.

116. A method as defined in claim 115 in which the weight vector generator step includes the step of generating the approximation for the limit point weight vector l_{LP} in accordance with the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2,$$

in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_2^2 + b_{i5}s_1s_2,$$

in an asymmetric case ("i" otherwise), in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

117. A method as defined in claim 116 in which weight vector generator step includes the step of selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

118. A method as defined in claim 114 in which the weight vector generator step includes the step of generating an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points

$\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

119. A method as defined in claim 118 in which the weight vector generator step includes the step of performing the extrapolation approximation methodology such that $M=3$.

120. A method as defined in claim 119 in which the weight vector generator step includes the step of generating the approximation for the limit point weight vector weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J),$$

where "J" is a predetermined integer and where

$$\left(l_{LP}(J)(s_1, s_2) \right)_m = \left(S_{sc, T, K, L, LP}(J)(s_1, s_2) \right)_{1, m},$$

where

$$S_{sc, T, K, L, LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc, T, K, L}(s_1(j), s_2(j))$$

with

$$S_{sc, T, K, L, LP}(0)(s_1, s_2) := I_{K+1}$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and

where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

121. A method as defined in claim 9 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the feature representation generator step includes the step of determining the tangent vector $e_c(q)$ in accordance with

$$e_c(q) = \sum_{i=0}^K (l_c(s_1, s_2))_i c^j(i)$$

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c = \lim_{j \rightarrow \infty} \frac{l_c(J)}{\|l_c(J)\|}$$

where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, and where

$$l_c(J) = (0, 1, 0, \dots, -1, 0, \dots) \cdot \prod_{j=J}^{J_p} S_{sc, T, K, L}(s_1(j), s_2(j))$$

where two non-zero components of the row vector on the right hand side are a "one" at position "one" in the row vector, and a "negative one" at position $\frac{K}{2} + 1$, and where $S_{sc, T, K, L}(s_1(j), s_2(j))$ corresponds to $S_{sc, T, K, L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product.

122. A method as defined in claim 121 in which the weight vector generator step includes the step of generating the tangent vector weight vector l_c using a polynomial approximation methodology.

123. A method as defined in claim 122 in which the weight vector generator step includes the step of generating the approximation for the tangent vector weight vector l_c in accordance with the polynomial

$$(l_c)_i = b_{i0}(s_1 + s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2) ,$$

in the anti-symmetric case (), or the polynomial

$$(l_c)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3 ,$$

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

124. A method as defined in claim 123 in which weight vector generator step includes the step of selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right) ,$$

where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

125. A method as defined in claim 121 in which the weight vector generator step includes the step of generating an approximation for the tangent vector weight vector l_c using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points

$\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0,...,M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

126. A method as defined in claim 125 in which the weight vector generator step includes the step of performing the extrapolation approximation methodology such that $M=3$.

127. A method as defined in claim 126 in which the weight vector generator step includes the step of generating the approximation for the tangent vector weight vector l_C in accordance with

$$l_C \approx \sum_{J=0}^3 b_J l_C(J)$$

where "J" is a predetermined integer and where

$$l_C(J)(s_1, s_2) = d(K)^J v_C \cdot S_{sc,T,K,L,LP}(J)(s_1, s_2)$$

where vector v_C is given by

$$v_C = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right)$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}}$$

and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j))$$

with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1}$$

where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

At
Cmt

128. A method as defined in claim 9 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being across the smooth feature line, the feature representation generator step includes the step of determining the tangent vector $e_c(q)$ in accordance with

$$e_s(q) = \sum_{i=0}^K (l_s(s_1, s_2))_i c^j(i)$$

6 where $l_s(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s = \lim_{j \rightarrow \infty} \frac{l_s(J)}{\|l_s(J)\|}$$

7

8 where $\|v\| = \sqrt{\sum_i v_i^2}$, that is, the Euclidean norm, where

100662152-020102

9

10

11

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares methodology and the values of the parameters s_1 and s_2 are at selected values.

131. A method as defined in claim 130 in which weight vector generator step includes the step of selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

132. A method as defined in claim 127 in which the weight vector generator step includes the step of generating an approximation for the tangent vector weight vector l_s using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points

$\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

133. A method as defined in claim 132 in which the weight vector generator step includes the step of performing the extrapolation approximation methodology such that $M=3$.

134. A method as defined in claim 133 in which the weight vector generator step includes the step of generating the approximation for the limit point weight vector weight vector in accordance with

$$l_s \approx \sum_{J=0}^3 b_J l_s(J),$$

4 where "J" is a predetermined integer and where

$$I_S(J)(s_1, s_2) = d(K)^J v_S \cdot S_{sc,T,K,L,LP}(J)(s_1, s_2),$$

5
6 where vector v_S is given by

$$v_S = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right),$$

7
8 dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}},$$

9
10 and

$$S_{sc,T,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,T,K,L}(s_1(j), s_2(j)),$$

11
12 with

$$S_{sc,T,K,L,LP}(0)(s_1, s_2) := I_{K+1},$$

13
14 where I_{K+1} is the "K+1" by "K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

135. A method as defined in claim 108 in which the mesh comprises a quadrilateral mesh in which, at a selected level "j," vertex $v_q(0)$ is at position $c^j(0)$ and neighboring points $v_q(k)$, $k=1, \dots, 2K$ are at respective positions $c^j(k)$, and in which the weight vector generator step includes the step of making use of a parameterized subdivision rule $S_{sc,Q,K,L}$ that relates the position $c^{j+1}(0)$ of the vertex $v_q(0)$ and positions $c^{j+1}(k)$ of neighboring points $v_q(k)$ at the next higher level "j+1" as follows

$$c^{j+1} = S_{sc,T,K,L} c^j$$

where subdivision rule $S_{sc,Q,K,L}$ is given by

10062192-020102

A1
Cmt

$$(S_{sc,Q,K,L}(s_1, s_2))_{l,m} = \begin{cases} (1-s_3)\left(1-\frac{7}{4K}\right) + \frac{3}{4}s_3 & \text{if } l=0, m=0 \\ (1-s_3)\left(\frac{3}{2K^2}\right) + \frac{1}{8}s_3 & \text{if } l=0, m=1 \text{ or } L+1 \\ (1-s_3)\left(\frac{3}{2K^2}\right) & \text{if } l=0, m=2, \dots, L \\ & \text{or } l=0, m=L+2, \dots, K \\ (1-s_3)\left(\frac{1}{4K^2}\right) & \text{if } l=0, m=K+1, \dots, 2K \\ \frac{3}{8}(1-s_2) + \frac{1}{2}s_2 & \text{if } l=1, m=0 \text{ or } 1 \\ \frac{1}{16}(1-s_2) & \text{if } l=L+1, m=2, K, \\ & K+1 \text{ or } 2K \\ \frac{3}{8}(1-s_1) + \frac{1}{2}s_1 & \text{if } l=L+1, m=0 \text{ or } L+1 \\ \frac{1}{16}(1-s_1) & \text{if } l=L+1, m=L, L+2, \\ & K+1 \text{ or } K+L+1 \\ \frac{3}{8} & \text{if } l=2, \dots, L, m=0 \\ & \text{or } l=L+2, \dots, K, m=0 \\ & \text{or } l=m=2, \dots, L \\ & \text{or } l=m=L+2, \dots, K \\ \frac{1}{16} & \text{if } l=2, \dots, L, m=l-1 \\ & \text{or } l=2, \dots, L, m=l+1 \\ & \text{or } l=2, \dots, L, m=K+l-1 \\ & \text{or } l=2, \dots, L, m=K+l \\ & \text{or } l=L+2, \dots, K, m=l-1 \\ & \text{or } l=L+2, \dots, K-1, m=l+1 \\ & \text{or } l=K, m=1 \\ & \text{or } l=L+2, \dots, K, m=K+l-1 \\ & \text{or } l=L+2, \dots, K, m=K+l \\ \frac{1}{4} & \text{if } l=K+1, \dots, 2K-1, m=0, \\ & l-K, l-K+1 \text{ or } l \\ & \text{or } l=2K, m=0, K, 1 \text{ or } 2K \\ 0 & \text{otherwise} \end{cases}$$

where the smooth feature line is defined in connection with the edges between respective points $v_q(1)$ and $v_q(L+1)$ ($L+1 \leq 2K$) and vertex $v_q(0)$, s_1 is the parameter associated with the edge between point $v_q(1)$ and vertex $v_q(0)$, s_2 is the parameter associated with the edge between point $v_q(L+1)$ and vertex $v_q(0)$, and $s_3 = \frac{1}{2}(s_1 + s_2)$.

136. A method as defined in claim 135 in which the representation of the feature is defined by at least one limit point associated with the vertex, the feature representation generator step includes the step of determining a position $\sigma(q)$ of the limit point in accordance with

$$\sigma(q) = \sum_{i=0}^{2K} (l_{LP}(s_1, s_2))_i c^j(i),$$

where $l_{LP}(s_1, s_2)$ is a vector of limit point weight values defined by

$$l_{LP}(s_1, s_2) = v_{LP} \cdot S_{sc, Q, K, L, LP}(s_1, s_2),$$

where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)),$$

where $S_{sc, Q, K, L}(s_1(j), s_2(j))$ corresponds to $S_{sc, Q, K, L}$ for sharpness parameters corresponding to the "j-th" level in the matrix product, where $S_{sc, Q, K, L, LP}(s_1, s_2)$ arguments s_1 and s_2 on the left-hand side refer to the sharpness parameters a definition level of the smooth feature line and the subscript "LP" refers to "Limit Point," and

$$v_{LP} = \frac{1}{K(K+5)}(K^2, 4, \dots, 4, 1, \dots, 1)$$

13

1 137. A method as defined in claim 136 in which the weight vector generator step includes the step
2 of generating an approximation for the limit point weight vector l_{LP} using a polynomial
3 approximation methodology.

1 138. A method as defined in claim 137 in which the weight vector generator step includes the step
2 of generating the approximation for the limit point weight vector l_{LP} in accordance with the
3 polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}(s_1 + s_2) + b_{i2}(s_1^2 + s_2^2) + b_{i3}s_1s_2, \quad ,$$

4 *Am*
5 in a symmetric case, or the polynomial

$$(l_{LP})_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2, \quad ,$$

6
7 in an asymmetric case, in which the coefficients b_{ij} are determined by a least squares methodology
8 and the values of the parameters s_1 and s_2 are at selected values.

1 139. A method as defined in claim 138 in which weight vector generator step includes the step of
2 selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right), \quad ,$$

3

4 where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

140. A method as defined in claim 134 in which the weight vector generator step includes the step of generating an approximation for the limit point weight vector l_{LP} using an extrapolation approximation methodology in relation to an M-degree polynomial that interpolates the points $\{x = 2^{-2^J}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the point $x=0$.

141. A method as defined in claim 140 in which the weight vector generator step includes the step of performing the extrapolation approximation methodology such that $M=3$.

142. A method as defined in claim 141 in which the weight vector generator step includes the step of generating the approximation for the limit point weight vector weight vector in accordance with

$$l_{LP} \approx \sum_{J=0}^3 b_J l_{LP}(J),$$

where "J" is a predetermined integer and where

$$(l_{LP}(J)(s_1, s_2))_m = (S_{sc, Q, K, L, LP}(J)(s_1, s_2))_{1, m},$$

where

$$S_{sc, Q, K, L, LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc, Q, K, L}(s_1(j), s_2(j)),$$

with

$$S_{sc, Q, K, L, LP}(0)(s_1, s_2) := I_{2K+1},$$

10 where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

143. A method as defined in claim 135 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being along the smooth feature line, the feature representation generator step includes the step of determining the tangent vector $e_c(q)$ in accordance with

$$e_c(q) = \sum_{i=0}^{2K} (l_c(s_1, s_2))_i c^j(i) ,$$

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_c(s_1, s_2) = d(K)^K v_c \cdot S_{sc, Q, K, L, LP}(s_1(j), s_2(j)) ,$$

where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

9
10 and vector v_c is defined as

$$(v_c)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \cos \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \cos \frac{2\pi(i-K-1)}{K} + \cos \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases},$$

where

$$A_K = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)},$$

and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_K}{16} + \frac{1}{4}}$$

144. A method as defined in claim 143 in which the weight vector generator step includes the step of generating the tangent vector weight vector l_c using a polynomial approximation methodology.

145. A method as defined in claim 144 in which the weight vector generator step includes the step of generating the approximation for the tangent vector weight vector l_c in accordance with the polynomial

$$(l_c)_i \approx b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2)$$

5 in the anti-symmetric case, or the polynomial

$$(l_c)_i \approx b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3,$$

6

7 in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares
8 methodology and the values of the parameters s_1 and s_2 are at selected values.

1 146. A method as defined in claim 145 in which weight vector generator step includes the step of
2 selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

3 where "N" is a selected integer, and indices $i, j = 0, \dots, N-1$.

1 147. A method as defined in claim 143 in which the weight vector generator step includes the step
2 of generating an approximation for the tangent vector weight vector l_c using an extrapolation
3 approximation methodology in relation to an M-degree polynomial that interpolates the points
4 $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the
5 point $x=0$.

1 148. A method as defined in claim 147 in which the weight vector generator step includes the step
2 of performing the extrapolation approximation methodology such that $M=3$.

1 149. A method as defined in claim 148 in which the weight vector generator step includes the step
2 of generating the approximation for the tangent vector weight vector l_c in accordance with

$$l_C \approx \sum_{J=0}^3 b_J l_C(J)$$

where "J" is a predetermined integer and where

$$l_C(J)(s_1, s_2) = d(K)^J v_C \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2)$$

where vector v_C is given by

$$v_C = \left(0, \cos \frac{2\pi(0)}{K}, \cos \frac{2\pi(1)}{K}, \dots, \cos \frac{2\pi(K-1)}{K} \right)$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}}$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) = \prod_{j=J}^{J_0} S_{sc,Q,K,L}(s_1(j), s_2(j))$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1}$$

where I_{K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

150. A method as defined in claim 135 in which the representation of the feature is defined by a tangent vector associated with the vertex, the tangent vector being across the smooth feature line, the feature representation generator step includes the step of determining the tangent vector $e_c(q)$ in accordance with

$$e_s(q) = \sum_{i=0}^{2K} (l_s(s_1, s_2))_i c^j(i) ,$$

where $l_c(s_1, s_2)$ is a vector of tangent vector weight values defined by

$$l_s(s_1, s_2) = d(K)^K v_s \cdot S_{sc, Q, K, L, LP}(s_1(j), s_2(j)) ,$$

where

$$S_{sc, Q, K, L, LP}(s_1, s_2) = \prod_{j=\infty}^{j_D} S_{sc, Q, K, L}(s_1(j), s_2(j)) ,$$

10 and vector v_c is defined as

$$(v_s)_i = \begin{cases} 0 & \text{if } i = 0 \\ A_k \sin \frac{2\pi(i-1)}{K} & \text{if } i = 1, \dots, K \\ \sin \frac{2\pi(i-K-1)}{K} + \sin \frac{2\pi(i-K)}{K} & \text{if } i = K+1, \dots, 2K \end{cases},$$

11 where

$$A_k = 1 + \cos\left(\frac{2\pi}{K}\right) + \cos\left(\frac{\pi}{K}\right) \sqrt{2\left(9 + \cos\frac{2\pi}{K}\right)},$$

14 and where dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{A_k}{16} + \frac{1}{4}}.$$

1 151. A method as defined in claim 150 in which the weight vector generator step includes the step
2 of generating the tangent vector weight vector l_c using a polynomial approximation methodology.

1 152. A method as defined in claim 151 in which the weight vector generator step includes the step
2 of generating the approximation for the tangent vector weight vector l_s in accordance with the
3 polynomial

$$(l_s)_i = b_{i0}(s_1 - s_2) + b_{i1}(s_1^2 - s_2^2) + b_{i2}(s_1^3 - s_2^3) + b_{i3}(s_1^2 s_2 - s_1 s_2^2),$$

5 in the anti-symmetric case, or the polynomial

$$(l_s)_i = b_{i0} + b_{i1}s_1 + b_{i2}s_2 + b_{i3}s_1^2 + b_{i4}s_1s_2 + b_{i5}s_2^2 + b_{i6}s_1^3 + b_{i7}s_1^2s_2 + b_{i8}s_1s_2^2 + b_{i9}s_2^3,$$

6

7 in the non-symmetric case, in which the coefficients b_{ij} are determined by a least squares
8 methodology and the values of the parameters s_1 and s_2 are at selected values.

1 153. A method as defined in claim 152 in which weight vector generator step includes the step of
2 selecting values of s_1 and s_2 in accordance with

$$(s_1, s_2) = \left(\cos \left(\frac{\left(i + \frac{1}{2} \right) \pi}{N} \right), \cos \left(\frac{\left(j + \frac{1}{2} \right) \pi}{N} \right) \right),$$

3
4 where "N" is a selected integer, and indices $i, j=0, \dots, N-1$.

1 154. A method as defined in claim 150 in which the weight vector generator step includes the step
2 of generating an approximation for the tangent vector weight vector l_s using an extrapolation
3 approximation methodology in relation to an M-degree polynomial that interpolates the points
4 $\{x = 2^{-2^j}, y = l_{LP}(J)\}$, $J=0, \dots, M$ and then evaluating this polynomial by extrapolation at the
5 point $x=0$.

1 155. A method as defined in claim 154 in which the weight vector generator step includes the step
2 of performing the extrapolation approximation methodology such that $M=3$.

1 156. A method as defined in claim 155 in which the weight vector generator step includes the step
2 of generating the approximation for the limit point weight vector weight vector in accordance with

$$l_S \approx \sum_{J=0}^3 b_J l_S(J)$$

where "J" is a predetermined integer and where

$$l_S(J)(s_1, s_2) = d(K)^J v_S \cdot S_{sc,Q,K,L,LP}(J)(s_1, s_2)$$

where vector v_S is given by

$$v_S = \left(0, \sin \frac{2\pi(0)}{K}, \sin \frac{2\pi(1)}{K}, \dots, \sin \frac{2\pi(K-1)}{K} \right)$$

dilation factor $d(K)$ is given by

$$d(K) = \frac{1}{\frac{3}{8} + \frac{1}{4} \cos \frac{2\pi}{K}}$$

and

$$S_{sc,Q,K,L,LP}(J)(s_1, s_2) := \prod_{j=J}^0 S_{sc,Q,K,L}(s_1(j), s_2(j))$$

with

$$S_{sc,Q,K,L,LP}(0)(s_1, s_2) := I_{2K+1}$$

where I_{2K+1} is the "2K+1" by "2K+1" identity matrix, and where

$$b_0 = \frac{-135}{120015}$$

$$b_1 = \frac{1270}{120015}$$

$$b_2 = \frac{-12192}{120015}$$

$$b_3 = \frac{131072}{120015}$$

Al
Coal

10062192-020102